

Early Stage Colon Cancer

Effective Date: May 2026



Background

In 2022, in Canada, 24300 people will develop colorectal cancer (13500 males and 10800 females).¹

A patient may be predisposed to develop colorectal cancer by a hereditary condition (e.g.: hereditary non-polyposis colon cancer, familial adenomatous polyposis) or a personal history of either inflammatory bowel disease (e.g.: Crohn's disease, ulcerative colitis) or adenomatous polyps. Over 60 percent of colorectal cancers are without a clearly identifiable predisposing factor, however.

In the absence of hereditary cancer syndromes, the progression from adenoma to adenocarcinoma occurs sporadically as a result of acquired genetic alterations. Allelic loss, chromosomal amplifications, or translocations account for 85 percent of such cases. In the other 15 percent, epigenetic silencing of a component of the DNA mismatch repair system allows frame-shift mutations and base-pair substitutions to persist. The resultant accumulation of tandemly repeated nucleotide sequences ("microsatellites") facilitate further changes during replication ("genetic instability"). While most microsatellites fail to occur within regulatory genes, microsatellites within critical coding regions of genes involved in the regulation of cell growth predispose to loss of function of a tumor suppressor genes or to the gain of function of an oncogene.

Patients with tumours that display "high-frequency microsatellite instability" (MSI-high) achieve a better five-year overall survival than patients with tumours that display microsatellite stability or low-frequency instability (HR 0.31, CI_{95%} 0.14-0.72, $p = 0.004$).² The use of adjuvant 5-Fluorouracil and Leucovorin chemotherapy fails to improve overall survival in patients with tumours that display high-frequency microsatellite instability (HR 1.07, CI_{95%} 0.62-1.86, $p = 0.80$).

After a diagnosis of colorectal cancer, prognosis depends upon the stage at diagnosis; that is, prognosis is better with less penetration of tumour into the bowel wall, fewer involved regional lymph nodes, and no evidence of metastatic disease.

Because the prognosis is better when colorectal cancer is identified at an earlier stage, because of the relatively high incidence of colorectal cancer, and because of the simplicity and accuracy of screening tests, screening for colorectal cancer represents an important component of routine care for all adults aged fifty years or older. This is especially important in patients with first-degree relatives with colorectal cancer.

Guideline Questions

1. What are the recommendations for the diagnostic workup of adult patients with potentially resectable colon cancer?
2. What are the recommendations for adjuvant chemotherapy in adult patients with colon cancer resected with curative intent and without evidence of metastatic disease?

Search Strategy

This guideline was developed to promote evidence-based practice in Alberta. It was compiled from the results of randomized controlled trials and systematic reviews, derived from an English language and relevant term search of PubMed and MEDLINE from 1990 forward. It takes into consideration related information presented at local, national, and international meetings as well as the Alberta Provincial Gastrointestinal Tumour Team's interpretation of the data. For the 2023 update of this guideline, recommendations were modified based on a consensus discussion at the 2023 Annual Gastrointestinal Tumour Team Meeting. However, no formal update of the literature was performed.

Target Population

The recommendations outlined in this guideline apply to adults over the age of 18 years with early stage colon cancer. Different principles may apply to pediatric patients.

Recommendations and Discussion

Suggested Diagnostic Work-Up

- Prior to an attempt at resection of an intraluminal mass, it is recommended that, in the absence of a complete bowel obstruction, a colonoscopy be completed to exclude synchronous neoplasms.
- A CT scan of the thorax, abdomen, and pelvis is recommended to exclude the possibility of metastatic disease and to provide a baseline for the surveillance CT scans. To evaluate an abnormality identified on CT scan, further imaging (e.g.: MR, ultrasound) may be required.
- A pre-operative CEA is recommended for future comparison. A post-operative CEA should be requested to ensure that it has normalized if it was elevated before surgery. ASCO recommends tumour budding should be assessed pre-operatively as well, however, at this time we believe the evidence is not strong enough to recommend adjuvant chemotherapy solely based on high levels of tumour budding.³
- The number of risk factors should be considered as part of the shared decision-making process. The presence of more than one risk factor may increase the risk of recurrence.⁴

Stage Information

Table 1. AJCC Cancer Staging System for Early Stage Colon Cancer, 8th Edition.

Stage	Depth of Tumour Penetration		Regional Lymph Node Involvement		Metastases	
Stage 0	T _{is}	Carcinoma <i>in situ</i>	N ₀	None	M ₀	Absent
Stage I	T ₁	Invasion into submucosa	N ₀	None	M ₀	Absent
	T ₂	Invasion into muscularis propria	N ₀	None	M ₀	Absent
Stage II _A	T ₃	Invasion through muscularis propria into peri-colic tissues	N ₀	None	M ₀	Absent
Stage II _B	T _{4a}	Tumor invades through the visceral peritoneum	N ₀	None	M ₀	Absent

Stage	Depth of Tumour Penetration		Regional Lymph Node Involvement		Metastases	
Stage II _c	T _{4b}	Direct invasion into, or adherence [§] to, other structures	N ₀	None	M ₀	Absent
Stage III _A	T ₁₋₂	As described above	N ₁	One to three lymph nodes	M ₀	Absent
			N _{1c}	Tumour deposits in subserosa, mesentery, or peri-rectal tissues		
	T ₁	Invasion into submucosa	N _{2a}	Four to six lymph nodes	M ₀	Absent
Stage III _B	T _{3-4a}	As described above	N ₁	One to three lymph nodes	M ₀	Absent
			N _{1c}	As described above		
	T ₂₋₃	As described above	N _{2a}	As described above	M ₀	Absent
	T ₁₋₂	As described above	N _{2b}	Seven or more lymph nodes	M ₀	Absent
Stage III _C	T _{4a}	Penetration to surface of visceral peritoneum	N _{2a}	As described above	M ₀	Absent
	T _{3-4a}	As described above	N _{2b}	As described above	M ₀	Absent
	T _{4b}	Direct invasion into, or adherence [§] to, other structures	N ₁₋₂	As described above	M ₀	Absent
Stage IV _A	T _{any}	As described above	N _{any}	As described above	M _{1a}	One site*
Stage IV _B	T _{any}	As described above	N _{any}	As described above	M _{1b}	Multiple sites*
Stage IV _C	T _{any}	As described above	N _{any}	As described above	M _{1c}	Metastasis involving peritoneal surface

* M_{1a} refers to metastasis confined to one site (e.g.: liver, lung, ovary, non-regional lymph node) whereas M_{1b} refers to metastasis in more than one site or within the peritoneum excluding the peritoneal surface.

§ If the microscopic assessment identifies no tumor in the adhesion, classify based on anatomic depth of wall invasion.

Note: A peri-tumoural nodule in the peri-colic adipose tissue without histologic evidence of lymph node architecture may represent discontinuous spread, venous invasion with extravascular spread, or a totally replaced lymph node.

Goal of Therapy

To render the patient free of disease and to delay or prevent recurrence.

Recommendations

1. After surgical resection of the tumour, the patient's need for adjuvant chemotherapy should be defined based upon the stage and the medical oncologist's assessment of the patient's relevant comorbidities (e.g.: diabetes mellitus, hypertension, cardiac status, etc.). The presence of at least one high-risk feature should prompt consideration for systemic adjuvant chemotherapy in stage II disease.
2. All patients should be tested for mismatch repair deficiency or microsatellite instability. If tumor testing is suspicious for a hereditary disposition to colorectal cancer or the patient presents with numerous polyps suggestive of an adenomatous polyposis syndrome, referral to the Genetics Counseling Service should be discussed with the patient.
3. Ideally, adjuvant chemotherapy should be initiated as soon as possible (once the patient has recovered from surgery).⁵⁻⁷ If this is not possible due to post-operative complications, adjuvant

chemotherapy could still be considered up to twelve weeks after surgery.

4. Consider treatment on a clinical trial, if available.
5. DPYD genotyping should be included in the planning of fluoropyrimidine based treatments [\[Link\]](#).
6. For stage II and III colon cancer, molecular profiling is recommended to determine somatic PI3K pathway alterations (group A: prespecified *PIK3CA* hotspot mutations in exon 9 or 20, or group B: moderate- or high-impact somatic variants in *PIK3CA*, *PIK3R1*, or *PTEN*). If somatic PI3K alterations are present, start aspirin (unless contra-indicated) 160 mg daily x 3 years after recovery (concurrent with chemotherapy if applicable)⁸. [Level of Evidence I]ⁱ
7. After completion of adjuvant chemotherapy, patients should be referred to a structured exercise program (Level of Evidence I).⁹
8. To optimize the care of patients with resected colon cancer, complicated cases should be reviewed with the multidisciplinary team.

Table 2. Recommendations for Adjuvant Treatment of Early Stage Colon Cancer.

Stage	Recommendations
Stage 0	- Although stage 0 disease is typically identified incidentally at the time of a colonoscopic polypectomy, inadequate margins warrants further endoscopic or surgical resection with oncologic principles. - No adjuvant systemic therapy is indicated.
Stage I	- Perform surgical resection with oncologic principles.^{10,11} - No adjuvant systemic therapy is indicated.
Stage II	- Perform surgical resection with oncologic principles.^{12,13} - Adjuvant chemotherapy should not routinely be offered to patients who are at low risk for recurrence, including:¹⁴⁻¹⁷ [Level of Evidence: IV; Strength of Recommendation: C] - Stage IIA (T3) tumours with at least 12 sampled lymph nodes - Absence of perineural or lymphovascular invasion, poor or undifferentiated tumour grade, clinical intestinal obstruction, tumour perforation - Less than grade BD3 tumour budding - If a “high-risk” or “poor prognosis feature” is present, consider adjuvant chemotherapy as for stage III disease. However, the benefit of Oxaliplatin in stage II disease has been questioned.¹⁸⁻²¹ “High-risk” features include direct invasion into adjacent structures, perforation through the tumor, clinical obstruction at presentation, poorly-differentiated histology, lymphovascular and/or perineural invasion, or evaluation of less than twelve regional lymph nodes. - There is no compelling evidence to suggest that age of patient should alter this recommendation. Specifically, there is no evidence that younger low-risk stage II patients should be offered ACT on the basis of their age alone. - Patients with proficient mismatch repair/microsatellite stable (pMMR or MSS) tumours are included within the previously mentioned recommendations (see “high risk features above”).

ⁱ In the ALASCAA trial, patients with stage I-III rectal cancer and stage II to III colon cancer were randomized to ASA or placebo within 12 weeks of surgery. The estimated 3-year cumulative incidence of recurrence was reduced with aspirin [group A: 7.7% versus 14.1% with placebo, HR 0.49 (95% CI 0.24 to 0.98), P=0.04; group B: 7.7% versus 16.8% with placebo, HR 0.42 (95% CI, 0.21 to 0.83)]. Severe adverse events occurred in 16.8% of aspirin recipients and 11.6% of placebo recipients. Testing and the cost of ASA are not currently funded.

Stage	Recommendations						
	- Systematic reviews have suggested that high levels of microsatellite instability confer a better overall survival as well as a possible resistance to 5-Fluorouracil.^{2,22,23} - For patients with mismatch repair deficiency (dMMR) or high microsatellite instability (MSI) tumours: - Adjuvant fluoropyrimidine-only chemotherapy is not routinely recommended [Level of Evidence: IV; Strength of Recommendation: C] - Poor differentiation is not considered a high-risk prognostic factor in patients with dMMR or MSI-high tumours - For patients with dMMR or MSI-high and T4 tumours and/or other high-risk features (with the exception of poor differentiation), oxaliplatin-containing chemotherapy may be considered. - Circulating tumor DNA (ctDNA) was identified as an emerging potential predictive factor; however, there is insufficient evidence regarding the predictive value of chemotherapy						
Stage III	- Perform surgical resection with oncologic principles.^{12,13} [REF] - When compared to no adjuvant chemotherapy therapy, 5-Fluorouracil-based regimens reduce the relative risk of death by 30 to 36%.²³⁻²⁵ - Adjuvant chemotherapy options include: <table border="1" data-bbox="277 884 1479 1703"> <tr> <td data-bbox="277 884 516 1234">CAPOX/ XELOX²⁶</td><td data-bbox="516 884 1479 1234">Refer to “Capecitabine: A Guide for Patient Care.” NO16968 demonstrated an improvement in both the disease-free survival (66.1% <i>versus</i> 59.8%, HR 0.80, CI_{95%} 0.69-0.93, <i>p</i> = 0.0045) and relapse-free survival (67.8% <i>versus</i> 60.9%, HR 0.78, CI_{95%} 0.67-0.92, <i>p</i> = 0.0024) when compared 5-Fluorouracil and Leucovorin by either the Mayo or Roswell Park regimens. Although overall survival was not statistically significantly superior (77.6% <i>versus</i> 74.2%, HR 0.87, CI_{95%} 0.72-1.05, <i>p</i> = 0.1486), the Kaplan-Meier curves continue to separate.</td></tr> <tr> <td data-bbox="277 1234 516 1585">Modified FOLFOX^{18,27}</td><td data-bbox="516 1234 1479 1585">This regimen requires placement of a central venous catheter (CVC), peripherally inserted central catheter (PICC line), or port. For the subset of patients with stage III disease, the MOSAIC trial suggested that the addition of Oxaliplatin to 5-Fluorouracil and Leucovorin improves five-year disease-free (66.4% <i>versus</i> 58.9%, HR 0.78, CI_{95%} 0.65-0.93, <i>p</i> = 0.005) and six-year overall survival (72.9% <i>versus</i> 68.7%, HR 0.80, CI_{95%} 0.65-0.97, <i>p</i> = 0.023). The benefit of Oxaliplatin in patients over seventy years of age has been questioned.^{10-12,20}</td></tr> <tr> <td data-bbox="277 1585 516 1703">Capecitabine²⁸</td><td data-bbox="516 1585 1479 1703">Refer to “Capecitabine: A Guide for Patient Care.” Capecitabine has been shown to be equally efficacious as, but less toxic and less resource intensive than, the “Mayo Regimen” (see below).</td></tr> </table> The optimal duration of Oxaliplatin-based adjuvant chemotherapy (CAPOX or FOLFOX) is 6 months. Based on the primary analysis of the IDEA collaboration, the non-inferiority of 3 months is not proven (HR1.07 95% CI 1.00-1.15) using a non-inferiority margin 1.12.²⁹ However, given the cumulative-dose dependent toxicity of oxaliplatin,³⁰ a duration of 3 months may be considered in select patients after an informed discussion about the	CAPOX/ XELOX ²⁶	Refer to “Capecitabine: A Guide for Patient Care.” NO16968 demonstrated an improvement in both the disease-free survival (66.1% <i>versus</i> 59.8%, HR 0.80, CI _{95%} 0.69-0.93, <i>p</i> = 0.0045) and relapse-free survival (67.8% <i>versus</i> 60.9%, HR 0.78, CI _{95%} 0.67-0.92, <i>p</i> = 0.0024) when compared 5-Fluorouracil and Leucovorin by either the Mayo or Roswell Park regimens. Although overall survival was not statistically significantly superior (77.6% <i>versus</i> 74.2%, HR 0.87, CI _{95%} 0.72-1.05, <i>p</i> = 0.1486), the Kaplan-Meier curves continue to separate.	Modified FOLFOX ^{18,27}	This regimen requires placement of a central venous catheter (CVC), peripherally inserted central catheter (PICC line), or port. For the subset of patients with stage III disease, the MOSAIC trial suggested that the addition of Oxaliplatin to 5-Fluorouracil and Leucovorin improves five-year disease-free (66.4% <i>versus</i> 58.9%, HR 0.78, CI _{95%} 0.65-0.93, <i>p</i> = 0.005) and six-year overall survival (72.9% <i>versus</i> 68.7%, HR 0.80, CI _{95%} 0.65-0.97, <i>p</i> = 0.023). The benefit of Oxaliplatin in patients over seventy years of age has been questioned. ^{10-12,20}	Capecitabine ²⁸	Refer to “Capecitabine: A Guide for Patient Care.” Capecitabine has been shown to be equally efficacious as, but less toxic and less resource intensive than, the “Mayo Regimen” (see below).
CAPOX/ XELOX ²⁶	Refer to “Capecitabine: A Guide for Patient Care.” NO16968 demonstrated an improvement in both the disease-free survival (66.1% <i>versus</i> 59.8%, HR 0.80, CI _{95%} 0.69-0.93, <i>p</i> = 0.0045) and relapse-free survival (67.8% <i>versus</i> 60.9%, HR 0.78, CI _{95%} 0.67-0.92, <i>p</i> = 0.0024) when compared 5-Fluorouracil and Leucovorin by either the Mayo or Roswell Park regimens. Although overall survival was not statistically significantly superior (77.6% <i>versus</i> 74.2%, HR 0.87, CI _{95%} 0.72-1.05, <i>p</i> = 0.1486), the Kaplan-Meier curves continue to separate.						
Modified FOLFOX ^{18,27}	This regimen requires placement of a central venous catheter (CVC), peripherally inserted central catheter (PICC line), or port. For the subset of patients with stage III disease, the MOSAIC trial suggested that the addition of Oxaliplatin to 5-Fluorouracil and Leucovorin improves five-year disease-free (66.4% <i>versus</i> 58.9%, HR 0.78, CI _{95%} 0.65-0.93, <i>p</i> = 0.005) and six-year overall survival (72.9% <i>versus</i> 68.7%, HR 0.80, CI _{95%} 0.65-0.97, <i>p</i> = 0.023). The benefit of Oxaliplatin in patients over seventy years of age has been questioned. ^{10-12,20}						
Capecitabine ²⁸	Refer to “Capecitabine: A Guide for Patient Care.” Capecitabine has been shown to be equally efficacious as, but less toxic and less resource intensive than, the “Mayo Regimen” (see below).						

Stage	Recommendations
	relative risks and benefits (Shi, JCO, 2017). Based on sub-group analyses from the IDEA collaboration, both regimen and risk group can be considered, though not randomized (regimen and risk group) or pre-planned (risk group) (Shi, JCO, 2017). 1. Patients with low risk stage III disease (T1-3,N1) who are fit enough to tolerate CAPOX, do not have inferior outcomes with 3 compared to 6 months of adjuvant chemotherapy (HR 0.85 95% CI 0.1-1.01). Not all patients may be eligible for CAPOX and if FOLFOX is a preferred regimen, 6 months remains standard (HR1.10, 95% CI 0.96-1.26). 2. For patients with high risk disease (T4 or N2), 6 months of either CAPOX (HR1.02, 95% CI 0.89-1.17) or FOLFOX remains standard. 3 months of FOLFOX in particular was clearly inferior to 6 months and should not be offered (HR1.16, 95% CI1.06-1.26). Other regimens may be considered in specific clinical situations: 1. Mayo Regimen:^{31,32} Six four-week cycles where Leucovorin 20 mg/m² IV followed by 5-Fluorouracil 425 mg/m² IV are administered daily for five consecutive days. 2. Roswell Park Regimen: Four eight-week cycles where Leucovorin (500 mg/m² IV over two hours) followed by 5-Fluorouracil (500 mg/m² IV bolus) are administered once every week for six weeks. For patients with resected stage III dMMR/MSI-H colon cancer, atezolizumab + oxaliplatin-based chemotherapy is the preferred option for (Level of Evidence I). The ATOMIC³³ trial demonstrated that the addition of 1 year of atezolizumab to 6 months of FOLFOX chemotherapy improved disease-free survival HR 0.50 (95% CI 0.35-0.73, P<0.001) in this population. There was no significant difference in overall survival. Grade 3 or 4 adverse events occurred in 84.1% of the patients who received atezolizumab with mFOLFOX6 and in 71.9% of those who received mFOLFOX6 alone. The ATOMIC trial did not evaluate a shorter duration of chemotherapy. This is not currently funded.

5. As long as resection of either a metachronous polyp, colorectal cancer, or a metastasis to liver or lung remains appropriate, surveillance is recommended (see [Clinical Practice Guidelines for Colorectal Surveillance](#)).


References

1. Brenner D, Poirier A, Woods R, et al. Projected estimates of cancer in Canada in 2022. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne*. 05/02/2022 2022;194(17)doi:10.1503/cmaj.212097
2. Ribic CM, Sargent DJ, Moore MJ, et al. Tumor microsatellite-instability status as a predictor of benefit from fluorouracil-based adjuvant chemotherapy for colon cancer. *New England Journal of Medicine*. 2003;349(3):247-257.
3. Baxter N, Kennedy E, Bergsland E, et al. Adjuvant Therapy for Stage II Colon Cancer: ASCO Guideline Update. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 03/10/2022 2022;40(8)doi:10.1200/JCO.21.02538
4. van Rooijen K, Derksen J, Verkooijen H, Vink G, Koopman M. Translation of IDEA trial results into clinical practice: Analysis of the implementation of a new guideline for colon cancer. *International journal of cancer*. 10/15/2022 2022;151(8)doi:10.1002/ijc.34149
5. Biagi JJ, Raphael MJ, Mackillop WJ, Kong W, King WD, Booth CM. Association between time to initiation of adjuvant chemotherapy and survival in colorectal cancer: a systematic review and meta-analysis. *JAMA*. 2011;305(22):2335-2342. doi://dx.doi.org/10.1001/jama.2011.749
6. Yang Y, Lu Y, Tan H, et al. The optimal time of starting adjuvant chemotherapy after curative surgery in patients with colorectal cancer. *BMC Cancer*. May 9 2023;23(1):422. doi:10.1186/s12885-023-10863-w
7. Gao P, Huang XZ, Song YX, et al. Impact of timing of adjuvant chemotherapy on survival in stage III colon cancer: a population-based study. *BMC Cancer*. Mar 1 2018;18(1):234. doi:10.1186/s12885-018-4138-7
8. Martling A, Hed Myrberg I, Nilbert M, et al. Low-Dose Aspirin for PI3K-Altered Localized Colorectal Cancer. *N Engl J Med*. Sep 18 2025;393(11):1051-1064. doi:10.1056/NEJMoa2504650
9. Courneya KS, Vardy JL, O'Callaghan CJ, et al. Structured Exercise after Adjuvant Chemotherapy for Colon Cancer. *N Engl J Med*. Jul 3 2025;393(1):13-25. doi:10.1056/NEJMoa2502760
10. Muto T, Oya M. Recent advances in diagnosis and treatment of colorectal T1 carcinoma. *Diseases of the Colon & Rectum*. 2003;46(10 Suppl):89.
11. Wang HS, Liang WY, Lin TC, et al. Curative resection of T1 colorectal carcinoma: risk of lymph node metastasis and long-term prognosis. *Diseases of the Colon & Rectum*. 2005;48(6):1182-1192.
12. Thomas KK, Francescatti AB, Vreeland TJ, et al. Standardization of Colon Resection for Cancer: An Overview of the American College of Surgeons Commission on Cancer Standard 5.6. *Ann Surg Oncol*. © 2023. Society of Surgical Oncology.; 2024:6-9. vol. 1.
13. Le Voyer TE, Sigurdson ER, Hanlon AL, et al. Colon cancer survival is associated with increasing number of lymph nodes analyzed: a secondary survey of intergroup trial INT-0089. *J Clin Oncol*. Aug 1 2003;21(15):2912-9. doi:10.1200/jco.2003.05.062
14. Jonker D, Spithoff K, Maroun J, Gastrointestinal Cancer Disease Site Group of Cancer Care Ontario's Program in Evidence-based C. Adjuvant systemic chemotherapy for stage II and III colon cancer following complete resection. 2008;
15. Gill S, Loprinzi CL, Sargent DJ, et al. Pooled analysis of fluorouracil-based adjuvant therapy for stage II and III colon cancer: who benefits and by how much? *Journal of Clinical Oncology*. 2004;22(10):1797-1806.
16. Benson AB, 3rd, Schrag D, Somerfield MR, et al. American Society of Clinical Oncology recommendations on adjuvant chemotherapy for stage II colon cancer. *Journal of Clinical Oncology*. 2004;22(16):3408-3419.
17. O'Connor ES, Greenblatt DY, LoConte NK, et al. Adjuvant chemotherapy for stage II colon cancer with poor prognostic features. *Journal of Clinical Oncology*. 2011;29(25):3381-3388. doi://dx.doi.org/10.1200/JCO.2010.34.3426
18. Andre TBCM-BLNMTJHTTCZMCPBJT-FIdG. Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer. *New England Journal of Medicine*. 2004;350(23):2343-2351.
19. Yothers G, O'Connell MJ, Allegra CJ, et al. Oxaliplatin as adjuvant therapy for colon cancer: updated results of NSABP C-07 trial, including survival and subset analyses. *Journal of Clinical Oncology*. 2011;29(28):3768-3774. doi://dx.doi.org/10.1200/JCO.2011.36.4539
20. Tournigand C, Andre T, Bonnetain F, et al. Adjuvant therapy with fluorouracil and oxaliplatin in stage II and elderly patients (between ages 70 and 75 years) with colon cancer: subgroup analyses of the Multicenter International Study of Oxaliplatin, Fluorouracil, and Leucovorin in the Adjuvant Treatment of Colon Cancer trial. *Journal of Clinical Oncology*. 2012;30(27):3353-3360. doi://dx.doi.org/10.1200/JCO.2012.42.5645

21. McCleary NJ, Meyerhardt JA, Green E, et al. Impact of age on the efficacy of newer adjuvant therapies in patients with stage II/III colon cancer: findings from the ACCENT database. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2013;31(20):2600-2606. doi:10.1200/JCO.2013.49.6638; 10.1200/JCO.2013.49.6638
22. Popat S, Hubner R, Houlston RS. Systematic review of microsatellite instability and colorectal cancer prognosis. *Journal of Clinical Oncology*. 2005;23(3):609-618.
23. Sargent DJ, Marsoni S, Thibodeau SN, et al. Confirmation of deficient mismatch repair (dMMR) as a predictive marker for lack of benefit from 5-FU based chemotherapy in stage II and III colon cancer (CC): a pooled molecular reanalysis of randomized chemotherapy trials. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2008;26(15S):4008.
24. Dube S, Heyen F, Jenicek M. Adjuvant chemotherapy in colorectal carcinoma: results of a meta-analysis. *Diseases of the Colon & Rectum*. 1997;40(1):35-41.
25. Sargent DJ, Patiyil S, Yothers G, et al. End points for colon cancer adjuvant trials: observations and recommendations based on individual patient data from 20,898 patients enrolled onto 18 randomized trials from the ACCENT Group. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2007;25(29):4569-4574. doi:10.1200/JCO.2006.10.4323
26. Haller DG, Tabernero J, Maroun J, et al. Capecitabine plus oxaliplatin compared with fluorouracil and folinic acid as adjuvant therapy for stage III colon cancer. *Journal of Clinical Oncology*. 2011;29(11):1465-1471.
27. Kuebler JP, Wieand HS, O'Connell MJ, et al. Oxaliplatin combined with weekly bolus fluorouracil and leucovorin as surgical adjuvant chemotherapy for stage II and III colon cancer: results from NSABP C-07. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2007;25(16):2198-2204. doi:10.1200/JCO.2006.08.2974
28. Twelves C, Wong A, Nowacki MP, et al. Capecitabine as adjuvant treatment for stage III colon cancer. *New England Journal of Medicine*. 2005;352(26):2696-2704.
29. Shi QS, AF. Shields, AF, et al. Prospective pooled analysis of six phase III trials investigating duration of adjuvant (adjuv) oxaliplatin-based therapy (3 vs 6 months) for patients (pts) with stage III colon cancer (CC): The IDEA (International Duration Evaluation of Adjuvant chemotherapy) collaboration. *J Clin Oncol*. 2017;35(Suppl_18)
30. André T, Boni C, Navarro M, et al. Improved overall survival with oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment in stage II or III colon cancer in the MOSAIC trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 07/01/2009 2009;27(19)doi:10.1200/JCO.2008.20.6771
31. O'Connell MJ, Mailliard JA, Kahn MJ, et al. Controlled trial of fluorouracil and low-dose leucovorin given for 6 months as postoperative adjuvant therapy for colon cancer. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 1997;15(1):246-250.
32. Group QC. Comparison of fluorouracil with additional levamisole, higher-dose folinic acid, or both, as adjuvant chemotherapy for colorectal cancer: a randomised trial. QUASAR Collaborative Group. *Lancet*. 2000;355(9215):1588-1596.
33. Sinicrope FA, Ou FS, Arnold D, et al. Atezolizumab plus FOLFOX for Stage III Mismatch Repair-Deficient Colon Cancer. *N Engl J Med*. Mar 26 2026;394(12):1155-1166. doi:10.1056/NEJMoa2507874

Appendix A:

Figure 1. Observed survival rates for 28,491 cases with adenocarcinoma of colon. Data from SEER 1973-2005.



Appendix B: Chemotherapy Dosing and Schedule

Chemotherapy	Dose/Schedule
CAPOX/XELOX	Eight three-week cycles where Oxaliplatin 130 mg/m ² is administered IV Q21d and Capecitabine 1,000 mg/m ² is administered PO Q12h for fourteen days.
FOLFOX	Twelve two-week cycles where · Oxaliplatin (85 mg/m² IV over two hours) and Leucovorin (400 mg/m² IV over two to six hours) are administered concurrently, · Followed by 5-Fluorouracil (400 mg/m² IV bolus), · Followed by a continuous intravenous infusion of 5-Fluorouracil (2,400 mg/m² over forty-six hours).
Capecitabine	Eight three-week cycles where Capecitabine 1,250 mg/m ² is administered PO Q12h for fourteen days.

Development and Revision History

This guideline was developed by a multidisciplinary working group comprised of members from the Alberta Provincial GI Tumour Team, external participants identified by the Working Group Lead, and a methodologist from the Guideline Resource Unit. The draft guideline was externally reviewed and endorsed by members of the Alberta Provincial GI Tumour Team who were not involved in the guideline's development, including surgical oncologists, radiation oncologists, medical oncologists, gastroenterologists, nurses, pathologists, and pharmacists. A detailed description of the methodology followed during the guideline development process can be found in the [Guideline Resource Unit Handbook](#).

This guideline was originally developed in 2008.

Levels of Evidence

I	Evidence from at least one large randomized, controlled trial of good methodological quality (low potential for bias) or meta-analyses of well-conducted randomized trials without heterogeneity
II	Small randomized trials or large randomized trials with a suspicion of bias (lower methodological quality) or meta-analyses of such trials or of trials with demonstrated heterogeneity
III	Prospective cohort studies
IV	Retrospective cohort studies or case-control studies
V	Studies without control group, case reports, expert opinion

Strength of Recommendations

A	Strong evidence for efficacy with a substantial clinical benefit; strongly recommended
B	Strong or moderate evidence for efficacy but with a limited clinical benefit; generally recommended
C	Insufficient evidence for efficacy or benefit does not outweigh the risk or the disadvantages (adverse events, costs, etc.); optional
D	Moderate evidence against efficacy or for adverse outcome; generally not recommended
E	Strong evidence against efficacy or for adverse outcome; never recommended

Maintenance

A formal review of the guideline will be conducted in 2024. If critical new evidence is brought forward before that time, however, the guideline working group members will revise and update the document accordingly.

Abbreviations

AJCC, American Joint Committee on Cancer; CAPOX, capecitabine + oxaliplatin; CEA, carcinoembryonic antigen; CI, confidence interval; CT, computed tomography; CVC, central venous catheter; FOLFOX, oxaliplatin + leucovorin + 5-fluorouracil; HR, hazard ratio; IV, intravenous; MR, magnetic resonance; PET, positron emission tomography; PICC, peripherally inserted central catheter; PO, by mouth, orally; SEER, Surveillance, Epidemiology, and End Results database; TNM, tumour-node-metastasis; XELOX, oxaliplatin + capecitabine.

Disclaimer

The recommendations contained in this guideline are a consensus of the Alberta Provincial GI Tumour Team and are a synthesis of currently accepted approaches to management, derived from a review of relevant scientific literature. Clinicians applying these guidelines should, in consultation with the patient, use independent medical judgment in the context of individual clinical circumstances to direct care.

Copyright © (2026) Alberta Health Services

This copyright work is licensed under the [Creative Commons Attribution-NonCommercial-NoDerivative 4.0 International license](#). You are free to copy and distribute the work including in other media and formats for non-commercial purposes, as long as you attribute the work to Alberta Health Services, do not adapt the work, and abide by the other license terms. To view a copy of this license, see <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

The license does not apply to AHS trademarks, logos or content for which Alberta Health Services is not the copyright owner.

Funding Source

Financial support for the development of Cancer Care Alberta's evidence-based clinical practice guidelines and supporting materials comes from the Cancer Care Alberta operating budget; no outside commercial funding was received to support the development of this document.

All cancer drugs described in the guidelines are funded in accordance with the Outpatient Cancer Drug Benefit Program, at no charge, to eligible residents of Alberta, unless otherwise explicitly stated. For a complete list of funded drugs, specific indications, and approved prescribers, please refer to the [Outpatient Cancer Drug Benefit Program Master List](#).

Conflict of Interest Statements

*Dr. Patricia Tang has nothing to disclose.

Derek Tilley has nothing to disclose.

*Working group lead

Citation

Tang, P(lead). Tilley, D. Cancer Care Alberta, Alberta Health Services (2026). Clinical Practice Guideline on Early Stage Colon Cancer, Version 9. Accessed [Month, Year]. Available from: www.ahs.ca/guru